

Asiago eclipsing binaries program. III. V570 Per^{*,**}

L. Tomasella¹, U. Munari¹, S. Cassisi², A. Siviero¹, S. Dallaporta³, R. Sordo¹, and T. Zwitter⁴

¹ INAF Osservatorio Astronomico di Padova, Sede di Asiago, 36012 Asiago (VI), Italy

² INAF Osservatorio Astronomico di Collurania, Via M. Maggini, 64100 Teramo, Italy

³ Via Filzi 9, I-38034 Cembra (TN), Italy

⁴ University of Ljubljana, Department of Physics, Jadranska 19, 1000 Ljubljana, Slovenia

Received date; accepted date

Abstract. The orbit and physical parameters of the previously unsolved double-lined eclipsing binary V570 Per, discovered by the Hipparcos satellite, are derived using high resolution Echelle spectroscopy and B , V photoelectric photometry. The metallicity from χ^2 analysis of the spectra is $[M/H]=+0.02\pm0.03$, and reddening from interstellar NaI and KI absorption lines is $E_{B-V}=0.023\pm0.007$. V570 Per is a well detached system, with shallow eclipses due to low orbital inclination and no sign of chromospheric activity. The two components have masses of 1.449 ± 0.006 and 1.350 ± 0.006 $M_{\odot}$ and spectral types F3 and F5, respectively. They are both still within the Main Sequence band ($T_1=6842\pm25$ K, $T_2=6562\pm25$ K from χ^2 analysis, $R_1=1.523\pm0.030$, $R_2=1.388\pm0.019$ $R_{\odot}$ derived forcing the orbital solution to conform to the spectroscopic light ratio) and are dynamically relaxed to co-rotation with the orbital motion ($V_{\text{rot}} \sin i_{1,2}=40$ and 36 (± 1) km sec^{-1}). The distance to V570 Per obtained from the orbital solution is 123 ± 2 pc, in excellent agreement with the revised Hipparcos distance of 123 ± 11 pc. The observed properties of V570 Per components are compared to available families of stellar evolutionary tracks, and in particular to BaSTI models computed on purpose for exactly the observed masses and varied chemical compositions. This system is interesting since both components have their masses in the range where the efficiency of convective core overshooting has to decrease with the total mass as a consequence of the decreasing size of the convective core during the central H-burning stage. Our numerical simulations show that, in order to match all empirical constraints including also the spectroscopical measurements, a small but not null overshooting is required, with efficiencies of $\lambda_{OV}=0.14$ and 0.11 for the 1.449 and 1.350 $M_{\odot}$ components, respectively. This confirms the finding of Paper II on the similar system V505 Per. At the ≈ 0.8 Gyr age of the system, the element diffusion has reduced the surface metallicity of the models from the initial $[M/H]=+0.17$ to $[M/H]=+0.02$, in perfect agreement with the spectroscopically derived $[M/H]=+0.02\pm0.03$ value.

Key words. stars: fundamental parameters – binaries: spectroscopic – binaries: eclipsing – star: individual: V570 Per

1. Introduction

In the present series of papers we derive global physical parameters for a selection of double-lined eclipsing binaries (SB2 EBs) by means of Echelle high resolution, high S/N spectroscopy and B , V photoelectric photometry. Orbital solutions provide masses and radii, reddening is measured from intensity of interstellar absorption lines and atmospheric analysis supplies effective temperatures, surface gravities, metallicity and rotational velocities. The physical parameters we derive are of high quality (1% accuracy regime) and are used to constrain the input physics of theoretical stellar models. We compute tailored evolutionary stellar models for the exact masses and chemical

mixtures observed in the components of the binaries. This allows to better focus on the effect expected, for ex., from overshooting and element diffusion. Siviero et al. (2004) and Tomasella et al. (2007), hereafter Paper I and Paper II, outline details of the type of data and methods used throughout this program.

V570 Per (HD 19457, HIP 14673) is a nearby eclipsing binary of early F spectral type, whose variability, characterized by a 1.9 days period, was discovered by Hipparcos satellite. As it will be seen in the next sections, its components do not show intrinsic variability. They are well separated, perfectly round, slow rotating and well within their Roche lobes, which make them proper tests for stellar models. V570 Per, though seen projected toward the α Per (Melotte 20) young open cluster, is a foreground star not physically associated to the cluster. This can be deduced from the comparison between the

* based mainly on data obtained with Asiago 1.82 m telescope

** Table 1 available in electronic form only

Table 2. Measured radial velocities of V570 Per.

#	HJD	Phase	Star 1			Star 2			<S/N>
			RV _⊙	ε	O-C	RV _⊙	ε	O-C	
30716	153.4401	0.773	+135.7	1.0	0.10	- 98.2	0.6	0.04	121
30796	154.5557	0.360	- 65.8	0.6	-0.78	+117.5	0.5	0.40	112
30842	155.4359	0.823	+124.8	1.0	-0.20	- 87.6	0.5	-0.73	126
30903	156.4674	0.366	- 63.8	0.5	-1.58	+113.9	0.4	-0.20	121
31161	165.4203	0.075	- 28.4	0.5	0.54	+ 78.9	0.6	0.56	130
31219	166.4546	0.619	+100.7	0.5	0.40	- 59.7	0.3	0.66	143
31268	167.4285	0.132	- 61.2	0.8	0.06	+112.4	0.5	-0.60	128
31317	169.5041	0.224	- 89.8	0.6	-0.14	+142.6	1.2	-0.90	130
31456	197.4553	0.928	+ 74.3	0.3	1.66	- 29.6	0.8	1.06	137
31779	206.3355	0.599	+ 89.0	0.5	-0.23	- 47.5	0.9	0.97	133
31821	209.3144	0.166	- 75.5	0.9	0.17	+129.3	1.1	0.80	105
31967	217.4346	0.438	- 20.0	0.7	0.48	+ 69.6	0.5	0.33	110
32080	225.3691	0.612	+ 97.0	0.8	0.43	- 55.7	0.4	0.64	117
33297	480.5559	0.854	+114.4	0.6	1.10	- 74.2	0.8	0.11	110
33318	485.4339	0.420	- 30.8	0.2	1.31	+ 82.0	0.5	0.26	102
33378	505.4620	0.956	+ 52.9	0.7	-1.02	- 10.3	0.9	0.28	113
33431	507.2637	0.904	+ 87.3	0.5	-0.17	- 45.8	0.5	0.78	123
33439	507.3552	0.952	+ 56.1	0.4	-0.57	- 12.7	0.6	0.82	137
33706	533.4932	0.702	+132.2	0.8	0.50	- 93.5	0.5	0.49	105
33868	561.2357	0.296	- 86.3	0.8	0.15	+139.3	1.4	-0.80	90
33870	561.3171	0.339	- 74.5	0.8	-0.68	+125.2	1.0	-1.30	137
33932	564.2289	0.871	+105.2	0.6	-0.20	- 66.2	0.8	-0.37	122
33965	564.5032	0.015	+ 12.8	0.5	0.72	+ 35.3	1.7	0.98	111
34066	570.2604	0.044	- 6.5	0.5	0.80	+ 57.8	0.3	0.60	140
36011	894.3444	0.530	+ 44.4	0.6	0.23	- 0.8	0.3	-0.68	95
36072	895.4091	0.090	- 39.8	0.9	-1.53	+ 88.1	0.8	-0.25	113
36080	895.5091	0.143	- 66.9	0.7	-0.52	+119.0	0.7	0.50	137
36123	896.4234	0.624	+102.7	0.7	-0.20	- 64.2	0.8	-1.06	142
36125	896.4457	0.636	+107.8	0.7	-1.00	- 70.8	0.9	-1.35	139
36126	896.4658	0.646	+112.9	0.5	-0.40	- 75.6	0.9	-1.29	143
36128	896.4883	0.658	+116.7	1.0	-1.60	- 80.9	1.0	-1.26	145

distance and proper motion of the cluster ($d=183\pm7$ pc, $\mu_{\alpha}^* = +22.47\pm0.16$ mas yr⁻¹, $\mu_{\delta} = -25.99\pm0.17$ mas yr⁻¹, from van Leeuwen 1999) and of V570 Per ($d=117\pm14$ pc, $\mu_{\alpha}^* = +52.20\pm0.85$ mas yr⁻¹, $\mu_{\delta} = -41.58\pm0.79$ mas yr⁻¹, from Hipparcos Catalogue). The revised Hipparcos distance to V570 Per is $d=123\pm11$ pc (van Leeuwen 2007).

A preliminary photometric and spectroscopic study of V570 Per was presented by Munari et al. (2001, hereafter M01). It was derived during an evaluation of the performances expected from Gaia (an ESA Cornerstone mission) on eclipsing binaries and, as such, the accuracy and abundance of input observational data (Hipparcos/Tycho photometry and Gaia-like ground based spectra) were necessarily lower than in the present study. A new study of the binary based on a much improved set of observational data is therefore justified, especially if supported by brand new reddening and atmospheric analysis.

2. Photometric data

The photometric observations of V570 Per were obtained in B and V (standard Johnson filters) with a 28 cm Schmidt-Cassegrain telescope and an Optec SSP5 photometer. The diaphragm has a diameter of 77 arcsec. There are no stars in the aperture brighter than $V=15.5$ mag (contributing less than 0.001 mag to recorded photometry). The instrumentation already proved to be very accurate and reliable (cf. Paper I and Paper II) and thus

perfectly suited to deal with the low amplitude eclipses of V570 Per (cf. Fig. 1).

The comparison star is HD 19805 (HIP 14980, $B_T=8.108\pm0.015$, $V_T=7.973\pm0.012$, spectral type B9.5 V) and the check star is TYC 3315 308 1 ($B_T=9.919\pm0.029$, $V_T=9.567\pm0.032$). Both the comparison and check stars are close to V570 Per on the sky (the distances being ~ 42 and ~ 18 arcmin respectively) so the atmospheric corrections are rather small. All the observations were obtained at heights over the horizon in excess of 30 degrees. The comparison star was measured against the check star at least once every observing run. In all, 34 measurements of the magnitude difference comparison–check star were collected. It remained constant through the whole observing campaign within a standard deviation of 0.006 mag. This confirms Hipparcos findings that both the comparison and the check stars are not variable, and thus well suited to serve in the photometry of V570 Per. Following Bessell (2000) transformations from Tycho to Johnson photometric system, we adopted for the comparison star $B=8.073$ and $V=7.957$.

In all, 446 measurements in B and 465 in V were collected for V570 Per between 2000 and 2003. They are listed in Table 1 (available electronic only). Each photometric point is actually the mean of 10 consecutive and independent measurements (every one lasting 5 sec) and the typical error of the mean is 0.006 mag in B and 0.005 mag in V . All the observations are corrected for atmospheric extinction and color equation via nightly calibration on Landolt’s equatorial fields.

The light curves of V570 Per in each band as well as the ($B-V$) color are shown in Fig. 1. Spectroscopic and photometric observations are well distributed in phase. The eclipses are very shallow due to the low inclination of the orbit ($i=77^\circ$), and this is the main reason for the absence of color variations during the eclipses (both stars remain essentially visible throughout the whole eclipses). The mean brightness out of eclipses is $B=8.505$ and $V=8.053$ mag. The dispersion of the B and V measurements about their mean value out of the eclipses ($\sigma_B=0.007$, $\sigma_V=0.006$) is only marginally higher than the accuracy of a single measurement. Thus any intrinsic variability of the amplitude larger than 0.007 mag should be ruled out.

3. Spectroscopic observations

The spectra of V570 Per were obtained in 1999–2002 with the Echelle+CCD spectrograph on the 1.82 m telescope operated by INAF Osservatorio Astronomico di Padova atop Mt. Ekar (Asiago). The instrumentation and observing set-up exactly match those described in Paper I, to which we refer for details of the observing mode. Here we recall that the wavelength region covered is 4500–9480 Å with a resolving power $R \sim 20\,000$. A journal of the observations is given in Table 2, in which the first three columns list the spectrum number (from the Asiago Echelle log book), the heliocentric JD (-2451000) and the orbital phase, while the last column gives the S/N per pixel of the




Fig. 1. The observed B , V , $B-V$ and radial velocity curves of V570 Per. In the radial velocity panel, the open circles indicate the hotter and more massive (primary) star, while the filled circles pertain to the cooler and less massive (secondary) star. The orbital solution from Table 3 is over-plotted to the observed data.

recorded stellar continuum averaged over the whole measured wavelength range (4890–5690 Å). The other columns lists, for both components separately and in km sec^{-1} , the radial velocity, its error and the difference with the computed orbit. We obtained 31 spectra with exposure times ranging from 1200 to 1800 sec, which guarantee a good S/N ratio (cf. Table 2) while avoiding smearing due to the orbital motion (1500 sec correspond to less than 1% in orbital period).

As outlined in Paper II, small residual spectrograph flexures are removed by cross-correlating the rich telluric absorptions complexes at 5880–5940, 6275–6310, 6865–7050 and 7160–7330 Å with a synthetic telluric absorption spectrum. Night-sky lines ([OI] and OH) and city lines (HgI and NaI) are used to check the accuracy of the wavelength calibration. Their mean velocity was found to be $0.0 \pm 0.1 \text{ km sec}^{-1}$ on every spectrum.

3.1. Radial velocities

In deriving the radial velocities of V570 Per we followed strictly the method outlined in details in Paper I. In short, six adjacent Echelle orders, covering with no inter-order gaps the 4890–5690 Å wavelength range, are measured via two-dimensional cross-correlation technique (TODCOR),

based on Zucker & Mazeh (1994) algorithm. These six Echelle orders are chosen because they are densely populated by absorption lines, in particular FeI and MgI, that perform particularly well in terms of radial velocity. As templates we use synthetic spectra with the appropriate temperatures, surface gravities and rotational velocities, selected from the Munari et al. (2005, hereafter M05) synthetic spectral atlas computed at the same 20 000 resolving power as the Echelle scientific spectra. The high reliability of our TODCOR-based radial velocities has been subjected to extensive tests described in Paper II. Repeated here on V570 Per data, they performed equally well, and we can conclude that no *external* systematic error in excess of 0.25 km sec^{-1} affect our radial velocities.

The results of the radial velocity measurements are reported in Table 2. The mean error of radial velocities is 0.6 km sec^{-1} for star 1, and 0.7 km sec^{-1} for star 2, as estimated from comparison of the radial velocities obtained separately from each of the six Echelle orders analysed here.

3.2. Reddening

The measurement of reddening is a key step in the determination of the absolute temperature scale (and therefore




Fig. 2. The NaI doublet (5890, 5896 Å) region for V570 Per. The interstellar components are well separated from the stellar ones.

of the distance) of eclipsing binaries. In spite of its short distance, some reddening is expected to affect V570 Per given its low galactic latitude ($l=145.18$, $b=-8.19$).

Our spectra cover the interstellar NaI (5890 and 5896 Å) and KI (7665 and 7699 Å) doublets which are excellent estimators of the reddening as demonstrated by Munari & Zwitter (1997). They calibrated a tight relation linking the NaI D2 (5890 Å) and KI (7699 Å) equivalent widths with the E_{B-V} reddening. On spectra obtained at quadratures, lines from both components are un-blended with the interstellar ones, which can be therefore accurately measured. An example of such a spectrum is presented in Fig. 2. We derived an equivalent width of 0.08 ± 0.03 Å for NaI D2 (5890 Å), which corresponds to $E_{B-V} = 0.023 \pm 0.007$ mag. This is the value adopted in the rest of the paper. At such a low reddening, no detectable KI interstellar line is expected, as confirmed by our spectra.

As mentioned above, V570 Per is seen in the foreground of α Per cluster. The cluster extends over a wide area and suffers from a patchy distribution of the reddening, which amounts to $E_{B-V} = 0.09$ according to Crawford & Barnes (1974), and $E_{B-V} = 0.11$ following Prosser (1992). It is worth noticing that, according to Perry & Johnston (1982), HD 19665, which lies less than 1 arcmin from V570 Per, is affected by $E_{B-V} = 0.022$ reddening.

4. Orbital solution

The orbital modeling is performed with version WD98K93 (Milone et al. 1992) of the Wilson-Devinney code (Wilson & Devinney 1971, Wilson 1998), which incorporates modified stellar atmospheres. The linear limb

Table 3. Orbital solution for V570 Per. The set of parameters in column 2 refer to the unconstrained solution described in Sect. 4, those in column 3 to the luminosity-constrained solution of Sect. 6. Formal errors to the solutions are given. d_{orb} , d_{HIP1997} and d_{HIP2007} are the distances derived by our orbital analysis, the original Hipparcos parallax (ESA 1997) and its recently revised value (van Leeuwen 2007), respectively.

	unconstrained		luminosity constrained	
P (d)	1.9009382	± 0.0000008	1.9009382	± 0.0000008
T_0 (HJD)	2451895.2371	± 0.0002	2451895.2371	± 0.0002
a ($R_{\odot}$)	9.0929	± 0.013	9.0929	± 0.012
V_{γ} (km sec^{-1})	+22.81	± 0.03	+22.81	± 0.03
$q = \frac{M_2}{M_1}$	0.9319	± 0.0008	0.9319	± 0.0008
i (deg)	77.44	± 0.34	77.44	± 0.32
e	0		0	
T_1 (K)	6842	± 25	6842	± 25
$T_1 - T_2$ (K)	262	± 70	280	± 35
Ω_1	7.00	± 0.43	6.93	± 0.13
Ω_2	7.28	± 0.50	7.16	± 0.08
r_1 (R_1/a)	0.1655	± 0.0193	0.1675	± 0.0031
r_2 (R_2/a)	0.1498	± 0.0123	0.1526	± 0.0019
R_1 ($R_{\odot}$)	1.50	± 0.11	1.523	± 0.030
R_2 ($R_{\odot}$)	1.36	± 0.11	1.388	± 0.019
M_1 ($M_{\odot}$)	1.4487	± 0.0062	1.4487	± 0.0058
M_2 ($M_{\odot}$)	1.3500	± 0.0059	1.3500	± 0.0055
$\log g_1$ (cgs)	4.25	± 0.06	4.23	± 0.02
$\log g_2$ (cgs)	4.30	± 0.07	4.28	± 0.01
$M_{\text{bol},1}$	3.13	± 0.17	3.11	± 0.05
$M_{\text{bol},2}$	3.54	± 0.18	3.49	± 0.04
d_{orb} (pc)	121	± 10	123	± 2
d_{HIP1997} (pc)	117	± 14		
d_{HIP2007} (pc)	123	± 11		

darkening coefficients are taken from van Hamme (1993) for the appropriate T_{eff} , $\log g$, $[M/H]$. The temperature of the primary adopted in the orbital solution, $T_1 = 6842 \pm 25$ K, is that determined by atmospheric analysis (see Sect. 5 below). It is in close agreement with indications from spectral classification and de-reddened colors. Comparing the spectrum of V570 Per in the 8480–8740 Å wavelength region with the Gaia spectral atlas of Munari & Tomasella (1999, which was compiled from observations of MK standards obtained with the same instrument and resolution of the V570 Per spectra discussed in this paper), a spectral classification F3V+F5V is derived. The temperature for the F3V primary, according to Bertone et al. (2004) calibration, is 6845 K. The color out of eclipse is $B - V = +0.452$ (± 0.005), that corrected for $E_{B-V} = 0.023$ (± 0.007), provides $(B - V)_0 = +0.429$ (± 0.009). Comparing with Drilling & Landolt (2000) color calibration for main sequence stars, it corresponds to a spectral type F4V, i.e. the mean spectral type of the two components of the binary.

The Gaia-like orbital solution of M01 was adopted as the set of initial values for input parameters in the orbital modeling (period P , epoch T_0 , semi-major axis a , barycentric radial velocity V_{γ} , mass ratio $q = M_1/M_2$, in-




Fig. 3. Comparison between observed (thick line) and synthetic (thin line) V570 Per spectra over two sample wavelength regions dominated by MgI lines (top) and FeI lines (bottom). Spectra at orbital phases 0.34 and 0.66 (# 33870 and 36128 in Table 2, respectively) are shown, which allows full and reversed splitting of the absorption lines of the two components. In each panel the lower curves represent (not to scale with the main spectra but in correct proportion between them) the contribution of each component of the binary to the formation of the observed spectrum at the given phase. The markers connect the shifted wavelengths of the same MgI (top panels) and FeI (bottom panels) lines in the spectra of the two components of the binary.

clination i , eccentricity e , modified Kopal potentials $\Omega_{1,2}$ and L_1/L_2 ratio in each pass-band).

We computed the solution using mode 2 of the Wilson-Devinney code, which is appropriate for detached binaries with no constraints on the potentials. The adopted bolometric albedos and exponents in the bolometric gravity brightening law were set to $A_1=A_2=0.5$ and $g_1=g_2=0.3$, respectively, as appropriate for convective atmospheres (Wilson 1998). As in Paper II, full orbital solution runs were carried out also with logarithmic and square-root limb darkening laws, as well as with various combinations of $A_{1,2}$ and $g_{1,2}$ in the range $0.5 \leq A_{1,2} \leq 1.0$ and $0.3 \leq g_{1,2} \leq 1.0$ (A and g are expected to be of the order of unity for radiative envelopes). No improvement in the formal accuracy of the solution was obtained. No orbital parameter was found to vary, as a results of these extensive experiments, by more than its formal error listed in the final orbital solution of Table 3. The latter was obtain with the following parameters for the linear limb darkening law: $x_{bol,1}=0.607$, $x_{bol,2}=0.664$, $x_{V,1}=0.407$, $x_{V,2}=0.428$,

$x_{B,1}=0.504$, $x_{B,2}=0.525$. No evidence was found for multiple reflection effects or third light presence.

The orbital period of V570 Per is stable. Our photometry provides five epochs of minima (all of them with an uncertainty of ± 0.00012 days): primary eclipses on HJD=2451895.23713, 2451920.44464, 2452986.37566 and secondary ones on HJD=2451951.31480, 2452158.51707. The Hipparcos and Tycho Catalogues (ESA 1997) give 2448500.1520 ± 0.001 as the time of primary minimum, revised to 2448500.1655 ± 0.003 in M01 re-analysis in combination with radial velocities. All these minima are well fitted by the 1.9009382 ± 0.0000008 day period of the orbital solution in Table 3.

V570 Per light and radial velocity curves do not show signature of an eccentric orbit. The circularity of the orbit is confirmed by initial modeling runs during which eccentricity was allowed to vary and nevertheless it did not drift away from zero. After a few such trials e was set to zero for the rest of orbital modeling.

Table 4. Atmospheric parameters (and their error of the mean, cf. Sect. 5) of V570 Per from the χ^2 fit to synthetic spectra. They are compared to the results from orbital solution for the elements in common (ΔT_{eff} , $\log g$, $V_{\text{rot}} \sin i$).

	χ^2 fit to synth. spectra		orbital solution	
$T_{\text{eff},1}$ (K)	6842	± 25		
$T_{\text{eff},2}$ (K)	6562	± 25		
$\Delta T_{\text{eff}} = T_{\text{eff},1} - T_{\text{eff},2}$ (K)	280	± 35	262	± 70
$\log g_1$ (cgs)	4.25	± 0.13	4.25	± 0.05
$\log g_2$ (cgs)	4.25	± 0.13	4.30	± 0.05
$V_{\text{rot},1} \sin i$ (km sec $^{-1}$)	40	± 5	40	± 1
$V_{\text{rot},2} \sin i$ (km sec $^{-1}$)	40	± 5	36	± 1
[M/H]	+0.02	± 0.03		
$L_{V,1}/(L_{V,1}+L_{V,2})$	0.60	± 0.02	0.59	± 0.12
$L_{V,2}/(L_{V,1}+L_{V,2})$	0.40		0.41	

The orbital solution of Table 3 is over-plotted to the observational data in Fig. 1. Their r.m.s is 0.006 mag for both B and V photometry, 0.8 km sec $^{-1}$ for radial velocities. The formal accuracy is 0.41% and 0.44% on the mass of the primary and secondary, respectively, and 7.3% and 8.1% on their radii. The latter is inflated by the low orbital inclination of the system ($i=77.44^\circ \pm 0.34$), that produces grazing eclipses of low amplitude ($\Delta B_I=0.116$, $\Delta B_{II}=0.094$, $\Delta V_I=0.117$, $\Delta V_{II}=0.098$ mag).

As a test, separate orbital solutions were carried out combining in turn B or V photometry alone with the radial velocities. We converged to the same orbital parameters of Table 3, with larger formal errors, as expected. Furthermore, the orbital solution was carried out on B and V photometric data together, without including radial velocities in the fitting process, and retaining i , M_1 , M_2 from the final solution of Table 3. Again, the orbital solution converged - within the formal errors - to the same set of parameters given in Table 3, with $R_1=1.55 \pm 0.02 R_\odot$ (instead of 1.50 ± 0.11) and $R_2=1.33 \pm 0.02 R_\odot$ (instead of 1.36 ± 0.11).

To check the accuracy of the derived orbital solution, we compared the corresponding distance with the Hipparcos parallax. In calculating the distance, we adopted from Bessell, Castelli and Plez (1998) a bolometric magnitude $M_{\text{bol},\odot}=4.74$ for the Sun and, for the corresponding atmospheric parameters, a bolometric correction $BC=0.00$ for both V570 Per components. The resulting distance, for the $E_{B-V}=0.023$ color excess derived in sect 3.2, is $d=121 \pm 10$ pc, in agreement with the 123 ± 11 pc revised Hipparcos parallax for V570 Per (van Leeuwen 2007 and private communication in advance of publication).

Varying the adopted value for the solar bolometric magnitude (e.g. from $M_{\text{bol},\odot}=4.72$ reported by Strazys and Kuriliene 1981, to $M_{\text{bol},\odot}=4.74$ by Livingston 2000, and $M_{\text{bol},\odot}=4.83$ by Popper 1980) and the corresponding

bolometric correction for F5V stars ($BC=-0.02$, -0.08 and -0.03 , respectively), the distance derived from the orbital solution changes by ± 2 pc.

5. Atmospheric analysis

To derive the atmospheric parameters of both V570 Per components, we performed a χ^2 fit against the synthetic spectral atlas of M05. It covers the 2500–10500 Å wavelength range at the same resolving power of our spectra ($R = 20\,000$). The M05 synthetic spectra are calculated with the revised solar abundances by Grevesse & Sauval (1998) and the new opacity distribution functions (ODFs) of Castelli & Kurucz (2004), throughout the whole HR diagram for 12 different rotational velocities, T_{eff} ranging from 3500 to 47500 K, $\log g$ from 0.0 to 5.0, [M/H] from -2.5 to $+0.5$, two values of α -enhancement ($[\alpha/\text{Fe}]=0.0$ and $+0.4$), and three micro-turbulent velocities (1, 2, 4 km sec $^{-1}$). The χ^2 fitting was restricted to the higher S/N spectra obtained at both quadratures. We limited the fitting to the same 4890–5590 Å wavelength range used to derive the radial velocities.

In the χ^2 fitting procedure we fixed only the radial velocities of the two components, while T_{eff} , $\log g$, $V_{\text{rot}} \sin i$, [M/H], micro-turbulent velocity and brightness ratio for both components were treated as free parameters. The results are given in Table 4. The listed errors are the error of the mean computed on the results obtained separately on the six Echelle orders covering the 4890–5590 Å wavelength range. Table 4 also provides a comparison with parameters in common with the orbital solution, and it fosters strong confidence in the results of the atmospheric analysis. Figure 3 highlights the goodness of the match between observed and synthetic spectra by over-plotting sample wavelength intervals of them.

In a configuration such as that of V570 Per, with similar stars and partial eclipses that are relatively shallow, the information on the relative sizes of the stars is weak. This reflects in the appreciable uncertainties of the radii in Table 3. The brightness ratio of the two stars as determined by the χ^2 fit offers an important and independent check on the accuracy of the radii. Such comparison is provided in Table 4 as brightness ratio of the two stars. It is carried out in the V band because at its effective wavelength are centered the six contiguous echelle orders used in our atmospheric analysis. The brightness ratios derived by χ^2 fit and orbital solution are the same, well within the respective errors. Also the TODCOR algorithm converged on the same 0.60/0.40 brightness ratio when deriving the radial velocities of the two components.

The V570 Per components rotate in synchronicity with the orbital motion (as highlighted by the comparison between observed and expected $V_{\text{rot},1,2} \sin i$ in Table 4), have spherical shapes ($R_{\text{L1}} = R_{\text{pole}}$) and are widely separated ($R/a \sim 0.16$). Under these circumstances, no tidal-induced chromospheric activity is expected. This is confirmed by the absence of emission line cores or emission veiling in the CaII H & K blue doublet or CaII far-red triplet lines, as




Fig. 4. Comparison of evolutionary tracks with observed parameters for V570 Per components (filled circles with error bars). The metallicity, helium content and mass of the fitting tracks are given in each panel. Padova, Yonsei-Yale and Granada tracks are interpolated from published ones, while BaSTI tracks have been computed on purpose for this work for the exact masses of the two components (1.4487 and $1.3500 M_{\odot}$).

verified at all orbital phases by comparison with synthetic spectra.

6. Constraining the stellar radii

The shallow eclipses displayed by V570 Per cannot sharply constrain the stellar radii, as indicated by the $\sim 8\%$ error on the radii obtained in the unforced orbital solution of Sect. 4, which is listed in the second column of Table 3. However, a relevant constrain is provided by the luminosity ratio of the two components as determined by χ^2 analysis and TODCOR fitting (cf Sect. 5 and last two lines of Table 4). We have therefore re-run the orbital solution with the Wilson-Devinney code, operated this time in mode 0, which is specific to the case when the luminosities are externally supplied. We then rapidly converged to the luminosity-constrained orbital solution listed in the third column of Table 3. It confirms the unforced orbital solution of Sect. 4, with an overall decrease of formal errors and in particular of those on stellar radii




Fig. 5. Comparison of V570 Per components with BaSTI tracks computed - for the exact observed masses - with and without overshooting (see Sect. 6 for details).

and on the distance to the binary system. The radii improved from 1.50 ± 0.11 , $1.36 \pm 0.11 R_{\odot}$ to 1.523 ± 0.030 , $1.388 \pm 0.019 R_{\odot}$, and the distance to the system from 121 ± 10 to 123 ± 2 pc. The latter is now identical to the revised Hipparcos distance (123 ± 11 pc, van Leeuwen 2007).

The comparison of V570 Per with theoretical stellar models carried out in next Sect. 7 and in Figures 4-6 is based on this luminosity-constrained orbital solution.

7. Observations vs theory

As it has been extensively reviewed in Paper II, binary systems with well-measured masses, radii, effective temperatures and heavy elements abundance provide stringent tests for evolutionary stellar models. Many issues can be investigated by comparing empirical evidence for the binary stars and theoretical predictions such as the efficiency of super-adiabatic convection in low-mass stars, the efficiency of diffusive processes (atomic diffusion and radiative levitation) and the real extension of convective core overshooting during the central H-burning stage. For intermediate mass stars, i.e. stars more massive than about $1.2 M_{\odot}$, this latter issue is really a long-standing problem (see e.g. Cassisi 2004 and references therein). An additional, still unsettled issue is how convective core overshooting reduces in response to a decrease in stellar mass and its canonical convective core (see Woo & Demarque 2001 for a detailed discussion).

In Paper II, we have briefly outlined the main results that have been obtained in this context by using empirical data for galactic Open Clusters of suitable age. We noticed the strong limitations imposed by the uncertainties in the distance, reddening and heavy elements




Fig. 6. Comparison between the observed radii for the V570 Per components (dashed lines), their uncertainty (error bars) and the BaSTI evolutionary tracks (solid lines) computed with overshooting and for exactly the observed individual masses. The resulting age for V570 Per is $\approx 790 \pm 60$ Myr.

abundances adopted for the clusters (cf. Vandenberg et al. 2007). A firmer contribution can be provided by eclipsing binary systems of suitable masses, with examples provided by V459 Cas (Sandberg Lacy et al. 2004), TZ For (Vandenberg et al. 2006), AI Phe (Andersen et al. 1988, Pietrinferni et al. 2004, hereafter P04) and V505 Per (Paper II). The above derived $1.449 \pm 0.006 M_{\odot}$ and $1.350 \pm 0.006 M_{\odot}$ masses for the two components of V570 Per qualify this binary too as a suitable constrain to stellar theoretical models and worth a detailed discussion.

V570 Per measured properties have been compared with the following sets of theoretical stellar models: BaSTI (P04, see also Cordier et al. 2007), Padova (Girardi et al. 2000), Yonsei-Yale (Yi et al. 2001, second release), and Granada (Claret et al. 2003), which have all been computed by assuming a scaled-solar heavy elements distribution.

The loci of theoretical isochrones and of V570 Per components on the HR diagram are shown in Fig. 4. For Yonsei-Yale, Granada and Padova models, we interpolated linearly both in mass and metallicity between the closest published isochrones. BaSTI models, on the other hand, have been computed on purpose for this paper, for exactly the measured masses of the two V570 Per components and exploring a fine grid in metal abundance, helium content, element diffusion, and amount of core overshooting (including canonical models with no overshooting). The overshooting was treated in the same way as in Paper II, and as extensively discussed by P04.

Padova and Granada models do not account for elements diffusion during the life of a star, and Yonsei-Yale does it only for helium. Thus, the metallicity of their fitting isochrones in Fig. 4 pertains to the initial composition when V570 Per was born, and not to its present

value at the stellar surface. It is therefore expected that the metallicities derived by fitting to Padova, Granada and Yonsei-Yale isochrones are *larger* than measured by spectroscopy. The best fitting BaSTI models have an initial metallicity $[M/H] = +0.17$ for both stars, that decreases to $[M/H] = +0.00$ and $+0.04$ for the 1.45 and $1.35 M_{\odot}$ components, respectively, by the time of V570 Per current 0.8 Gyr age (see below for its derivation). The average of this values very well compares with the spectroscopically measure of $[M/H] = +0.02 \pm 0.03$ for the V570 Per components.

The comparison with theoretical isochrones in Fig. 4 shows that both components of V570 Per are only marginally evolved. The more massive of the two stars has already moved slightly away from the zero age main sequence. Figure 5 shows the effect of overshooting, which has been computed for BaSTI models according to the same P04 prescriptions already adopted in Paper II. This figure shows that canonical stellar models with no overshooting but accounting for element diffusion, can still reproduce the position of V570 Per in the H-R diagram (dashed line in Fig. 5). However, the canonical models give a surface metallicity at the present age of the binary system equal to $[M/H] = +0.12$, which is outside the range of uncertainty for the observed $[M/H] = +0.02 \pm 0.03$ value. On the other hand, a small amount of overshooting allows a perfect fit for both the spectroscopic metallicity and the H-R diagram location, as illustrated by the solid lines in Fig. 5 that have been computed for an efficiency of the overshooting amounting to $\lambda_{OV} = 0.14$ for the $1.449 \pm 0.006 M_{\odot}$ primary and $\lambda_{OV} = 0.11$ for the $1.350 \pm 0.006 M_{\odot}$ secondary. In Paper II, for the lighter components of V505 Per, we obtained proportionally lower efficiencies: $\lambda_{OV} = 0.093$ and 0.087 for the 1.269 and $1.251 M_{\odot}$ components, respectively. The efficiency is usually defined in terms of the parameter λ_{OV} that gives the length - expressed as a fraction of the local pressure scale height H_P - crossed by the convective cells in the convective stable region outside the Schwarzschild convective boundary. Regardless of the initial metallicity, P04 adopt $\lambda_{OV} = 0.20 \times H_P$ for $M \geq 1.7 M_{\odot}$, $\lambda_{OV} = 0$ for $M \leq 1.1 M_{\odot}$, and $\lambda_{OV} = (\frac{M}{M_{\odot}} - 0.9)/4$ for $1.1 M_{\odot} \leq M \leq 1.7 M_{\odot}$.

To estimate the age of the system, in Fig. 6 we plot the evolution in radius of the BaSTI models above described for the two V570 Per components and compare them with the observed values. The derived system age is 790 ± 60 Myr.

Acknowledgements. We would like to thank the anonymous referee for valuable suggestions, R. Barbon and E. Nasi for useful discussion, F. van Leeuwen who kindly communicated the revised Hipparcos parallax of V570 Per in advance of publication, and the technical staff operating the 1.82m telescope in Asiago for the skillful assistance during the whole project.

References

- Andersen, J., Clausen, J.V., Gustafsson, B., Nordström, B., Vandenberg, D.A. 1988, A&A 196, 128

- Bertone, E., Buzzoni, A., Chavez, M., Rodriguez-Merino, L.H. 2004, AJ, 128, 829
- Bessell, M.S., Castelli F., Plez, B. 1998, A&A, 333, 231
- Bessell, M.S. 2000, PASP, 112, 961
- Cassisi, S. 2004, in *Variable Stars in the Local Group*, IAU Colloquium 193, Christchurch (New Zealand), D. W. Kurtz and K. R. Pollard eds., ASP Conference Proceedings, Vol. 310, 489
- Castelli, F., Kurucz, R.L., 2004, A&A, 419, 725
- Claret A., Paunzen E., Maitzen H.M. 2003, A&A, 412, 91
- Cordier, D., Pietrinferni, A., Cassisi, S., Salaris, M. 2007, AJ, 133, 468
- Crawford, D.L., Barnes J.V. 1974, AJ, 79, 687
- Drilling, J.S., Landolt, A.U. 2000 in Allen's Astrophysical Quantities, A.N.Cox ed., Springer, 381
- ESA 1997, *The Hipparcos and Tycho Catalogues*, ESA SP-1200
- Girardi, L., Bressan, A., Bertelli, G., Chiosi, C. 2000, A&AS 141, 371
- Grevesse, N., Sauval, A.J. 1998, Space Sci Rew, 85, 161
- Lang, K.R. 2006, in *Astrophysical Formulae*, 3rd edition, Springer
- Livingston, W.C. 2000 in Allen's Astrophysical Quantities, A.N.Cox ed., Springer, 339
- Milone, E.F., Stagg, C.R., Kurucz, R.L. 1992, ApJS 79, 123
- Munari, U., Zwitter, T. 1997, A&A, 318, 269
- Munari, U., Tomasella, L. 1999, A&AS, 137, 521
- Munari, U., Tomov, T., Zwitter, T., Milone, E.F., Kallrath, J., Marrese, P.M., Boschi, F., Prsa, A., Tomasella, L., Moro, D. 2001, A&A, 378, 477
- Munari, U., Sordo, R., Castelli, F., Zwitter, T. 2005, A&A, 442, 1127
- Perry, C.L., Johnston, L. 1982, ApJ, 50, 451
- Popper, D.M. 1980, ARA&A 18, 115
- Pietrinferni, A., Cassisi, S., Salaris, M., Castelli, F. 2004, ApJ, 612, 168
- Prosser, C.F. 1992, AJ, 103, 488
- Sandberg Lacy, C.H., Claret, A., Sabby, J.A. 2004, AJ, 128, 132
- Siviero, A., Munari, U., Sordo, R., Dallaporta, S., Marrese, P.M., Zwitter, T.; Milone, E.F. 2004, A&A, 417, 1083 (Paper I)
- Straizys, V., & Kuriliene, G. 1981, Ap&SS 80, 353
- Tomasella, L., Munari, U., Siviero, A., Cassisi, S., Dallaporta, S., Zwitter, T., Sordo, R. 2007, A&A in press, astro-ph/0712.3387 (Paper II)
- Vandenberg, D.A., Bergbush, P. A., Dowler, P.D. 2006, ApJS, 162, 375
- Vandenberg, D.A., Gustafsson, B., Edvardsson, B., Eriksson, K., Ferguson, J., 2007, ApJ Letter, in press
- van Leeuwen, F. 1999, A&A, 341, L71
- van Leeuwen, F. 2007, *Hipparcos, the new reduction*, Springer
- van Hamme, W. 1993, AJ, 106, 2096
- Wilson, R.E., Devinney, E.J. 1971, ApJ, 166, 605
- Wilson, R.E. 1998, *Computing Binary Star Observables*, Univ. of Florida Astronomy Dept.
- Woo, J.-H., Demarque, P. 2001, AJ, 122, 1602
- Yi, S., Demarque, P., Lejeune, T., Barnes, S. 2001, ApJS 136, 417 2nd edition, Cambridge University Press
- Zucker, S., Mazeh, T. 1994, ApJ, 420, 806